



YOUR OCREVUS[®] & OCREVUS ZUNOVO[™] WELCOME GUIDE

Use this step-by-step guide to help you get ready for treatment.

BRITTA

USING OCREVUS
SINCE 2018

*For OCREVUS, your first dose will be split into 2, for a total of 3 treatments your first year.

For patients with relapsing multiple sclerosis (RMS) or primary progressive multiple sclerosis (PPMS)

What are OCREVUS and OCREVUS ZUNOVO?

OCREVUS and OCREVUS ZUNOVO are prescription medicines used to treat:

- Relapsing forms of multiple sclerosis (MS), to include clinically isolated syndrome, relapsing-remitting disease, and active secondary progressive disease, in adults
- Primary progressive MS, in adults.

It is not known if OCREVUS and OCREVUS ZUNOVO are safe and effective in children.

Who should not receive OCREVUS or OCREVUS ZUNOVO?

Do not receive OCREVUS or OCREVUS ZUNOVO if you:

- have an active hepatitis B virus (HBV) infection.
- have had a life-threatening administration reaction to ocrelizumab.
- have had a life-threatening allergic reaction to ocrelizumab, hyaluronidase, or any of the ingredients of OCREVUS ZUNOVO. Tell your healthcare provider if you have had an allergic reaction to OCREVUS or OCREVUS ZUNOVO or any of their ingredients in the past.

Visit [OCREVUS.com](https://www.ocrevus.com) or click for [OCREVUS Prescribing Information and Medication Guide](#), and [OCREVUS ZUNOVO Prescribing Information and Medication Guide](#).

IMPORTANT SAFETY INFORMATION

What is the most important information I should know about OCREVUS and OCREVUS ZUNOVO?

OCREVUS and OCREVUS ZUNOVO can cause serious side effects, including:

- **Infusion reactions (OCREVUS):** Infusion reactions are a common side effect of OCREVUS, which can be serious and may require you to be hospitalized. You will be monitored during your infusion and for at least 1 hour after each infusion of OCREVUS for signs and symptoms of an infusion reaction.
- **Injection reactions (OCREVUS ZUNOVO):** Injection reactions are a common side effect of OCREVUS ZUNOVO, which can be serious and may require you to be hospitalized. You will be monitored for signs and symptoms of an injection reaction when you receive OCREVUS ZUNOVO. This will happen during all injections for at least 1 hour after your first injection, and for at least 15 minutes after all injections following the first injection.

OCREVUS[®]
ocrelizumab



OCREVUS ZUNOVO[™]
ocrelizumab & hyaluronidase-ocsq
Subcutaneous injection 920mg



HELLO!

Inside this Welcome Guide you'll find information about what to expect during your treatment with OCREVUS or OCREVUS ZUNOVO.

We hope you'll take some time to review this information and keep it handy throughout your treatment. It is important to remember that while this is a helpful resource, you should always talk with your healthcare provider about any questions you may have.



Would you like additional support? Call OCREVUS CONNECTS®

With **OCREVUS CONNECTS®**, you can get helpful support and resources for your treatment journey.

Just call **1-844-OCREVUS (627-3887)** to speak with someone who can answer your questions about OCREVUS, OCREVUS ZUNOVO, and MS or turn to page 10 to find out more about OCREVUS CONNECTS®.

WHAT'S INSIDE

Overview

- 4 About OCREVUS & OCREVUS ZUNOVO

Treatment Experience

- 6 OCREVUS Infusions
- 7 OCREVUS ZUNOVO Injections
- 8 Infusion & Injection Reactions

FAQs

- 9 FAQs

Support

- 10 OCREVUS CONNECTS®
- 11 Patient Navigators & Clinical Education Managers (CEMs)
- 12 Financial Assistance

Additional Resources


- 14 Treatment Appointment Checklist
- 15 Questions to Ask Your Healthcare Team
- 16 Suggestions for Support Partners
- 18 Indications and Important Safety Information

HOW 2-TIMES-A-YEAR OCREVUS IS GIVEN

OCREVUS is an **infusion therapy** — also known as IV therapy — that is given through an IV placed in your arm. OCREVUS ZUNOVO is a **subcutaneous injection** given under the skin in your abdomen area. Both OCREVUS and OCREVUS ZUNOVO are administered by a healthcare professional.

For OCREVUS infusions, your first dose will be 2 separate infusions 2 weeks apart, each lasting 2.5 hours.* After your first dose, you will receive 1 infusion every 6 months, each lasting 2-4 hours. For OCREVUS ZUNOVO, there is no split dose, and you will receive 1 injection every 6 months, each lasting about 10 minutes.*

OCREVUS YEAR 1: THREE INFUSIONS





OCREVUS YEAR 2 & BEYOND: TWO INFUSIONS PER YEAR



OCREVUS ZUNOVO YEAR 1 & BEYOND: TWO INJECTIONS PER YEAR



*This is only administration time and does not account for all aspects of treatment. Actual clinic time may vary.
The duration of an infusion or injection could be longer should a reaction occur.

2-TIMES-A-YEAR* OCREVUS FOR RMS & PPMS AVAILABLE TWO WAYS	
4 HOUR [†] / 2 HOUR ^{†‡}	~10 MIN [†]
INFUSION	INJECTION
 OCREVUS [®] ocrelizumab <small>INTRAVENOUS INJECTION 920mg</small>	 OCREVUS ZUNOVO [™] ocrelizumab & hyaluronidase-ocsq <small>Subcutaneous injection 920mg</small>
Intravenous infusion	Subcutaneous injection

The duration of an infusion or injection could be longer should a reaction occur

*For OCREVUS, your first dose will be split into 2, for a total of 3 treatments the first year.

[†]This is only administration time and does not account for all aspects of treatment. Actual clinic time may vary.

^{†‡}For eligible patients who have not previously experienced any serious infusion reactions with OCREVUS.

After talking to your doctor, you may be able to receive OCREVUS or OCREVUS ZUNOVO at their office, an infusion center, or at home from a home health provider. You can discuss where to get your treatment with your doctor, and your healthcare team can help you find a location.

At some treatment providers, you may be in a private room or a shared space. Each provider has its own policy about what you can or cannot bring with you, so you may want to reach out to your healthcare team prior to your appointment.



Important Safety Information for OCREVUS and OCREVUS ZUNOVO (cont.) OCREVUS and OCREVUS ZUNOVO can cause serious side effects, including:

• Infection:

- Infections are a common side effect. OCREVUS and OCREVUS ZUNOVO increase your risk of getting upper respiratory tract infections, lower respiratory tract infections, skin infections, and herpes infections. Serious infections can happen with OCREVUS and OCREVUS ZUNOVO, which can be life-threatening or cause death. Tell your healthcare provider right away if you have an infection. Your healthcare provider should delay your treatment with OCREVUS or OCREVUS ZUNOVO until your infection is gone.

Visit [OCREVUS.com](https://www.ocrevus.com) or click for [OCREVUS Prescribing Information and Medication Guide](#), and [OCREVUS ZUNOVO Prescribing Information and Medication Guide](#).



WHAT AN OCREVUS INFUSION APPOINTMENT MAY LOOK LIKE:



PRE-INFUSION: 30-60 MINUTES

- You will be given premedication 30-60 minutes before starting your OCREVUS treatment to help reduce possible infusion reactions. This premedication can be given as pills or as part of your IV infusion



INFUSION: 2-4 HOURS

- You'll be seated in a supportive chair. Once settled, a thin needle will be inserted into your arm to start the infusion of OCREVUS. An automatic pump will also be set up to ensure you receive the exact amount
- If an infusion reaction occurs, your healthcare team may stop or slow the rate of your infusion



POST-INFUSION: 1 HOUR

- You will be asked to stay for at least an hour for observation. This will allow your healthcare team to watch out for any side effects related to your treatment

TOTAL APPOINTMENT TIME: AT LEAST 3:30-6 HOURS



Hear from people like Pastell about their OCREVUS infusion experience.

WATCH NOW

YOU MAY BE ABLE TO GET A SHORTER 2-HOUR* INFUSION

If you don't experience any serious infusion reactions with the 4-hour OCREVUS infusion, your next infusion could take as little as 2 hours.† It's the same dose of OCREVUS, just given faster. **Ask your doctor if the shorter 2-hour infusion is right for you.**

*This is only administration time and does not account for all aspects of treatment. Actual clinic time may vary.

†The duration of an infusion could be longer should a reaction occur.

Important Safety Information for OCREVUS and OCREVUS ZUNOVO (cont.) OCREVUS and OCREVUS ZUNOVO can cause serious side effects, including:

- Hepatitis B virus (HBV) reactivation:** If you have ever had hepatitis B virus infection, the hepatitis B virus may become active again during or after treatment with OCREVUS or OCREVUS ZUNOVO. Hepatitis B virus becoming active again (called reactivation) may cause serious liver problems including liver failure or death.
- Vaccinations:** Tell your doctor about any recent or upcoming vaccinations.
 - You should receive any required 'live' or 'live-attenuated' vaccines at least 4 weeks before you start treatment with OCREVUS or OCREVUS ZUNOVO.
 - When possible, you should receive any 'non-live' vaccines at least 2 weeks before you start treatment with OCREVUS or OCREVUS ZUNOVO.

Visit [OCREVUS.com](https://www.ocrevus.com) or click for [OCREVUS Prescribing Information and Medication Guide](#), and [OCREVUS ZUNOVO Prescribing Information and Medication Guide](#).

OCREVUS®
ocrelizumab
200mg/100mL PREPARED IN WATER

OCREVUS ZUNOVO™
ocrelizumab & hyaluronidase-ocsq
Subcutaneous injection 920mg

WHAT AN OCREVUS ZUNOVO INJECTION APPOINTMENT MAY LOOK LIKE:

For OCREVUS ZUNOVO, there is no split first dose.



PRE-INJECTION: 30 MINUTES

- You will take premedication at least 30 minutes before starting each OCREVUS ZUNOVO treatment, usually as pills to swallow
- You may be able to take your medications at home—your healthcare team will advise on all pre-treatment protocols



INJECTION: ~10 MINUTES

- You will be asked to sit or lie down, and a thin needle will be inserted under the skin in your abdomen area. This needle will be connected to the prepared syringe containing OCREVUS ZUNOVO
- A member of your healthcare team will either inject OCREVUS ZUNOVO manually or will use an automatic pump to do this



POST-INJECTION: 15 MIN-1 HOUR*

- Your healthcare team will observe you for at least one hour after your first injection. For every injection after that, you will be observed for at least 15 minutes
- This will allow your healthcare team to watch out for any side effects related to your treatment

*Depending on whether it is the first dose or a subsequent dose.

TOTAL APPOINTMENT TIME: AT LEAST 55 MIN-1:40 MIN

**STILL HAVE QUESTIONS
ABOUT OCREVUS ZUNOVO?**

[CLICK HERE TO
WATCH VIDEO](#)

OCREVUS ZUNOVO™
ocrelizumab & hyaluronidase-ocsq
Subcutaneous injection 920mg



Important Safety Information for OCREVUS and OCREVUS ZUNOVO (cont.) OCREVUS and OCREVUS ZUNOVO can cause serious side effects, including:

- **Weakened immune system:** OCREVUS or OCREVUS ZUNOVO taken before or after other medicines that weaken the immune system could increase your risk of getting infections.
- **Progressive Multifocal Leukoencephalopathy (PML):** PML is a rare brain infection that usually leads to death or severe disability and has been reported with ocrelizumab. Symptoms of PML get worse over days to weeks.
- **Decreased immunoglobulins:** OCREVUS and OCREVUS ZUNOVO may cause a decrease in some types of immunoglobulins. Your healthcare provider will do blood tests to check your blood immunoglobulin levels.

Visit [OCREVUS.com](https://www.ocrevus.com) or click for [OCREVUS v](#) and [Medication Guide](#), and [OCREVUS ZUNOVO Prescribing Information](#) and [Medication Guide](#).

OCREVUS®
ocrelizumab
SUBCUTANEOUS INJECTION



OCREVUS ZUNOVO™
ocrelizumab & hyaluronidase-ocsq
Subcutaneous injection 920mg



INFUSION & INJECTION REACTIONS

Infusion (OCREVUS) and injection (OCREVUS ZUNOVO) reactions are a common side effect, and can be serious and require you to be hospitalized. You will be monitored during your treatment for signs and symptoms of a reaction. Reactions are most common during treatment and within an hour after but can occur up to 24 hours afterward.

Please tell your healthcare provider or nurse if you get any of these symptoms:

- itchy skin
- rash
- hives
- tiredness
- coughing or wheezing
- trouble breathing
- throat irritation or pain
- feeling faint
- fever
- redness on your face (flushing)
- nausea
- headache
- swelling of the throat
- dizziness
- shortness of breath
- fatigue
- fast heartbeat

Additionally, for OCREVUS ZUNOVO: Injection site pain, swelling, or redness.

In a clinical study, injection reactions were more frequently reported with first injection. All reactions were mild to moderate, and treatable with treatment adjustments and medicines to help with reactions.

My infusion nurse is great about monitoring me throughout the entire process. So, before every infusion, she gives me medicine to help manage any potential reactions and keeps me an hour afterward.

JUDI
USING OCREVUS SINCE 2017

Important Safety Information for OCREVUS and OCREVUS ZUNOVO (cont.)

What are the possible side effects of OCREVUS and OCREVUS ZUNOVO?

OCREVUS and OCREVUS ZUNOVO can cause serious side effects, including:

- **Risk of cancers (malignancies) including breast cancer:** Follow your healthcare provider's instructions about standard screening guidelines for breast cancer.
- **Inflammation of the colon, or colitis:** Tell your healthcare provider if you have any symptoms of colitis, such as diarrhea, blood in stool, and stomach pain.
- **The most common side effects of OCREVUS ZUNOVO include:** injection reactions, respiratory tract infections, and skin infections.

Visit [OCREVUS.com](https://www.OCREVUS.com) or click for [OCREVUS Prescribing Information and Medication Guide](#), and [OCREVUS ZUNOVO Prescribing Information and Medication Guide](#).



FAQs

Q: How can OCREVUS or OCREVUS ZUNOVO affect pregnancy?

A: There is limited data on the effect OCREVUS or OCREVUS ZUNOVO has on pregnancy, and it is unknown if these treatments will harm your unborn baby. You should use birth control (contraception) during treatment with OCREVUS or OCREVUS ZUNOVO and for 6 months after your last treatment. Tell your healthcare provider if you are pregnant, think you might be pregnant, or plan to become pregnant.

Q: Can I receive vaccines while taking OCREVUS or OCREVUS ZUNOVO?

A: Since you may receive some kinds of vaccines but not others, it's important to discuss the vaccine, the timing of getting it, and your OCREVUS or OCREVUS ZUNOVO treatment with your healthcare provider. With OCREVUS or OCREVUS ZUNOVO, it is not recommended to receive 'live-attenuated' or 'live' vaccines while on treatment.

Q: What happens if I have a schedule conflict with my treatment or miss a dose of OCREVUS or OCREVUS ZUNOVO?

A: If a planned treatment of OCREVUS or OCREVUS ZUNOVO is missed, it should be given as soon as possible. Do not wait until the next planned dose. Doses must be separated by at least 5 months, so if you have a schedule conflict, contact your treatment provider to reschedule.

Q: What is a subcutaneous injection?

A: A subcutaneous (SC, SQ, or Sub-Q) injection is given into the fatty tissue under the skin. OCREVUS ZUNOVO is a subcutaneous injection given by a healthcare provider.

Q: Who can receive OCREVUS ZUNOVO?

A: OCREVUS ZUNOVO was developed for people with MS who may benefit from the efficacy, safety, and 2-times-a-year dosing of OCREVUS but prefer a route of administration different from infusions. You and your doctor can decide if it's the right choice for you.

Q: Why isn't there a split first dose for OCREVUS ZUNOVO?

A: OCREVUS was developed with the first dose being split into two infusions to evaluate the risk of reactions in each patient by slowing the amount of medicine absorbed into the bloodstream. Because OCREVUS ZUNOVO is administered subcutaneously, the medicine is absorbed into the bloodstream more slowly than through an OCREVUS infusion, so a split dose is not required.

Q: Why are there different active doses of OCREVUS ZUNOVO and OCREVUS?

A: The administration of OCREVUS ZUNOVO is a single 920mg injection under the skin of the abdomen. This dose is higher than the 600mg OCREVUS intravenous dose because of the different routes to the bloodstream.

With intravenous medications: The drug is directly administered into the vein, so these medications directly reach the blood circulation.

With subcutaneous medications: The drug is administered into the subcutaneous tissue, which may impact the amount of medicine and/or the time it takes for the medicine to reach the bloodstream.

Therefore, a higher dose of OCREVUS ZUNOVO is needed to achieve a medical effect similar to OCREVUS.



We're here for you


OCREVUS CONNECTS® is a resource center dedicated to answering your questions related to OCREVUS, OCREVUS ZUNOVO, and MS.

We're here to help you with treatment resources, insurance coverage questions, and more. Once OCREVUS or OCREVUS ZUNOVO has been prescribed for you, you can choose to enroll in OCREVUS CONNECTS® where you may be teamed up with either a Patient Navigator or a Clinical Education Manager (CEM) or both.

To enroll, call **1-844-OCREVUS (627-3887)**. You'll speak directly with a Patient Navigator who will connect you with the services that will best meet your needs. Support is available Monday through Friday, 9 AM to 8 PM ET, in English and Spanish.

OCREVUS CONNECTS® is a resource to answer your questions about OCREVUS, OCREVUS ZUNOVO, and MS. But remember, you should talk with your healthcare provider if you have any questions about your treatment.

THE SUPPORT SERVICES WE OFFER



Patient Navigators*
your guide throughout
your OCREVUS
treatment journey



Clinical Education Managers†
for one-on-one
supplemental education
sessions



Additional resources
including answers to questions
about OCREVUS, OCREVUS
ZUNOVO, and MS

*If you have enrolled in the program.

†Clinical Education Managers are Genentech employees and do not give medical advice.

Visit [OCREVUS.com](https://www.ocrevus.com) or click for [OCREVUS Prescribing Information and Medication Guide](#), and [OCREVUS ZUNOVO Prescribing Information and Medication Guide](#).




Subcutaneous injection 920mg



SUPPORT ALONG YOUR JOURNEY WITH OCREVUS AND OCREVUS ZUNOVO

WHAT DOES A PATIENT NAVIGATOR DO?



Helps you understand your insurance coverage and financial assistance options for your treatment



Answers questions about infusions and injections, and reviews the treatment process with you from start to finish



Coordinates finding a treatment site or gives you information about getting treatment at home to discuss with your doctor



Works with you and your healthcare team throughout your entire treatment journey

WHAT DOES A CLINICAL EDUCATION MANAGER* DO?

*This program is intended primarily for patients prescribed OCREVUS and OCREVUS ZUNOVO who are in their first year of treatment. CEMs are Genentech employees and do not give medical advice.



Offers one-on-one supplemental education sessions to help you better understand OCREVUS, OCREVUS ZUNOVO, and MS



Helps you prepare for your first treatment by explaining what to expect



Details local MS resources that may be available in your community for you to further connect with



REACH OUT FOR SUPPORT TODAY

Enroll with OCREVUS CONNECTS® today.
Call [1-844-OCREVUS](tel:1-844-OCREVUS) (627-3887), 9 AM-8 PM ET.

Visit [OCREVUS.com](https://www.ocrevus.com) or click for [OCREVUS Prescribing Information and Medication Guide](#), and [OCREVUS ZUNOVO Prescribing Information and Medication Guide](#).

OCREVUS®
ocrelizumab 200mg/10mL INJECTION

OCREVUS ZUNOVO™
ocrelizumab & hyaluronidase-ocsq
Subcutaneous injection 920mg



COMMITTED TO HELPING YOU FIND FINANCIAL ASSISTANCE OPTIONS

We understand your medication can be costly. If you are worried about paying for your treatment, a Patient Navigator may be able to find assistance options to help you pay for OCREVUS or OCREVUS ZUNOVO.

There are options if you have:

- A commercial insurance plan (such as one you get through your employer or through a Health Insurance Marketplace like HealthCare.gov)
- A public insurance plan (such as Medicare or Medicaid)
- No insurance

If you are eligible, the options to help you pay for OCREVUS and OCREVUS ZUNOVO include:

- The OCREVUS Co-pay Program* (if you have a commercial insurance plan)
- Referrals to independent co-pay assistance foundations[†]
- Referrals to the Genentech Patient Foundation[‡]

To learn more about how a Patient Navigator can help, call **1-844-OCREVUS (627-3887)**.

*Eligibility criteria apply. Not valid for patients using federal or state government programs for their Genentech medicine. Patients must be taking the Genentech medicine for an FDA-approved indication. Please visit the [Co-pay Program website](#) for the full list of Terms and Conditions.

[†]Independent co-pay assistance foundations have their own rules for eligibility. Genentech has no involvement or influence in independent foundation decision-making or eligibility criteria and does not know if a foundation will be able to help you. We can only refer you to a foundation that supports your disease state. This information is provided as a resource for you. Genentech does not endorse or show preference for any particular foundation. The foundations in this list may not be the only ones that might be able to help you.

[‡]If you have health insurance, you should try to get other types of financial assistance, if available. You also need to meet income requirements. If you do not have insurance, or if your insurance does not cover your Genentech medicine, you must meet a different set of income requirements.

ADDITIONAL RESOURCES

As you navigate your journey with OCREVUS, OCREVUS ZUNOVO, and MS, there are a few resources that may help (noted on pages 13-16). You can fill out, print, or take them with you to your next appointment with your healthcare provider or share them with loved ones.

INCLUDED ARE:



An Appointment checklist
to help prepare



Questions to ask your
healthcare team or
treatment provider



Tips for
support partners

I realize I have a proactive healthcare team, but I still have to
advocate for myself, and you should too.

.....
KATRINA
USING OCREVUS SINCE 2017

PREPARING FOR YOUR TREATMENT

YOUR OCREVUS APPOINTMENT CHECKLIST

The better prepared you are, the better your treatment experience will be. Here are a few suggestions to help get ready for your OCREVUS or OCREVUS ZUNOVO appointment.

BEFORE YOUR TREATMENT

Take care of these things the week before your treatment.

- ☐ Check with your treatment provider to find out if:
 - You can bring food or snacks
 - You can have a friend with you
 - There will be other people in the same room getting treatment
 - You need to officially visit the center prior to your treatment appointment
 - There are any premedication protocols to be aware of
- ☐ Read the Medication Guide for the OCREVUS treatment you are prescribed
- ☐ Plan ahead to have someone pick you up if you think you'll want support
- ☐ Call OCREVUS CONNECTS® at **1-844-OCREVUS (627-3887)** to ask a Patient Navigator any questions you may still have



THINGS TO BRING

Collect these items before your treatment appointment so you're ready to go.

- ☐ Insurance card and prior authorization from your insurer if requested
- ☐ A list of your medications and healthcare providers
- ☐ Water, food, snacks or gum if your treatment provider's policy allows them
- ☐ Activities to help pass the time, like a laptop or tablet, books, magazines, games, or puzzles

THE DAY OF YOUR TREATMENT

Be comfortable — wear loose-fitting clothes so your healthcare team can easily reach your arm or abdomen. You can also wear additional layers or bring a blanket to help control your warmth.

- ☐ Drink plenty of water to stay hydrated
- ☐ Depending on the OCREVUS treatment you are prescribed, be prepared for other medications you will receive before your treatment that could make you sleepy

AFTER YOUR TREATMENT

Don't forget to check in with your treatment provider and your doctor.

- ☐ Book your next treatment before you leave
- ☐ Talk to your doctor about your options for your next treatment appointment



**SCAN THIS CODE to watch stories
from real people using OCREVUS.**


Visit [OCREVUS.com](https://www.ocrevus.com) or click for [OCREVUS Prescribing Information and Medication Guide](#), and [OCREVUS ZUNOVO Prescribing Information and Medication Guide](#).



OCREVUS ZUNOVO™
ocrelizumab & hyaluronidase-ocsq
Subcutaneous injection 920mg



QUESTIONS TO ASK YOUR HEALTHCARE TEAM ABOUT YOUR OCREVUS OR OCREVUS ZUNOVO TREATMENT



Q: Is treatment at home an option for me? _____

Q: Can I continue taking my regular medications? _____

Q: Can I eat and drink before and during my treatment? _____

Q: How will I feel during my treatment? _____

Q: Will I notice any changes in my MS afterward? _____

Q: How will I feel after my treatment? _____

Q: What side effects might I experience after receiving treatment? _____

Q: How should I prepare for my treatment? _____

Q: What should I do if I start to feel like I'm having a reaction to my treatment? _____

Q: Will I be in a room by myself or will I be with other people? _____

Q: Can my loved one stay in the room with me to keep me company? _____

Q: Will I be able to talk on the phone during treatment? _____

Q: Can I drive a car immediately after? _____

Q: Can I go back to work after my treatment? _____

Q: If I am receiving treatment at home, how will that change any of the above? _____

NOTES:

SUGGESTIONS FOR SUPPORT PARTNERS

As your loved one gets ready for their treatment with OCREVUS or OCREVUS ZUNOVO, there may be things you can do to help you both feel prepared. Here are some tips to consider that may be useful during each step of the process.



BEFORE THE TREATMENT:

- Make a copy of the list of medications your loved one is taking, just in case you need it, including prescription and over-the-counter medicines, vitamins, and herbal supplements
- Keep a list of all of your loved one's healthcare provider information handy — you never know what information may be required
- Review treatment materials ahead of time
- If this is your loved one's first treatment, they may be nervous — let them know you're available to talk
- Keep a reminder of their appointment somewhere visible, like on the refrigerator

DAY OF TREATMENT:

- Consider asking your loved one if they would like company during their treatment, whether at the treatment provider or their home
- Help them prepare for their appointment — have a bag ready filled with comfort items such as a blanket, warm socks, a laptop, or puzzles, and bring water to help them stay hydrated
- If you aren't able to accompany your loved one to their treatment, try to make sure you're available in case they want to reach you
- If you are staying during the treatment and have questions, a nurse should be able to provide information

SUGGESTIONS FOR SUPPORT PARTNERS (CONT.)



AFTER TREATMENT:

- Look for any signs of side effects — you may notice things that your loved one does not
- Note that reactions are most common during treatment and within an hour after but can occur up to 24 hours later
- Check in with them the day after their treatment to see how they're feeling
- Consider creating reminders — add scheduled treatment dates to your calendar so you'll know when they're coming up
- Try to be as flexible as possible with your loved one and how they may feel afterward
- Keep the conversation going — having regular touch points with your loved one can be helpful for both of you

Understand the OCREVUS and OCREVUS ZUNOVO treatment journey better.

Call **OCREVUS CONNECTS®** at **1-844-OCREVUS (627-3887)**
9 AM-8 PM ET if you have questions.

I took the leap with OCREVUS and I'm glad I did.

.....
HANNAH
USING OCREVUS SINCE 2018

INDICATIONS AND IMPORTANT SAFETY INFORMATION

What are OCREVUS and OCREVUS ZUNOVO?

OCREVUS and OCREVUS ZUNOVO are prescription medicines used to treat:

- Relapsing forms of multiple sclerosis (MS), to include clinically isolated syndrome, relapsing-remitting disease, and active secondary progressive disease, in adults
- Primary progressive MS, in adults.

It is not known if OCREVUS and OCREVUS ZUNOVO are safe and effective in children.

Who should not receive OCREVUS or OCREVUS ZUNOVO?

Do not receive OCREVUS or OCREVUS ZUNOVO if you:

- have an active hepatitis B virus (HBV) infection.
- have had a life-threatening administration reaction to ocrelizumab.
- have had a life-threatening allergic reaction to ocrelizumab, hyaluronidase, or any of the ingredients of OCREVUS ZUNOVO. Tell your healthcare provider if you have had an allergic reaction to OCREVUS or OCREVUS ZUNOVO or any of their ingredients in the past.

What is the most important information I should know about OCREVUS and OCREVUS ZUNOVO?

OCREVUS and OCREVUS ZUNOVO can cause serious side effects, including:

- **Infusion reactions (OCREVUS):** Infusion reactions are a common side effect of OCREVUS, which can be serious and may require you to be hospitalized. You will be monitored during your infusion and for at least 1 hour after each infusion of OCREVUS for signs and symptoms of an infusion reaction.
- **Injection reactions (OCREVUS ZUNOVO):** Injection reactions are a common side effect of OCREVUS ZUNOVO, which can be serious and may require you to be hospitalized. You will be monitored for signs and symptoms of an injection reaction when you receive OCREVUS ZUNOVO. This will happen during all injections for at least 1 hour after your first injection, and for at least 15 minutes after all injections following the first injection.

Tell your healthcare provider or nurse if you get any of these symptoms:

- | | | |
|------------------------|-----------------------------|--------------------------|
| ○ itchy skin | ○ throat irritation or pain | ○ headache |
| ○ rash | ○ feeling faint | ○ swelling of the throat |
| ○ hives | ○ fever | ○ dizziness |
| ○ tiredness | ○ redness on your face | ○ shortness of breath |
| ○ coughing or wheezing | (flushing) | ○ fatigue |
| ○ trouble breathing | ○ nausea | ○ fast heartbeat |

Additionally, for OCREVUS ZUNOVO:

- | | | |
|-----------------------|------------|-----------|
| ○ injection site pain | ○ swelling | ○ redness |
|-----------------------|------------|-----------|

These infusion and injection reactions can happen during or up to 24 hours after administration.

It is important that you call your healthcare provider right away if you get any of the signs or symptoms listed above after each infusion or injection.

INDICATIONS AND IMPORTANT SAFETY INFORMATION (CONT.)

- **Infection:**

- o Infections are a common side effect. OCREVUS and OCREVUS ZUNOVO increase your risk of getting upper respiratory tract infections, lower respiratory tract infections, skin infections, and herpes infections. Serious infections can happen with OCREVUS and OCREVUS ZUNOVO, which can be life-threatening or cause death. Tell your healthcare provider if you have an infection or have any of the following signs of infection including fever, chills, or a cough that does not go away, or painful urination. Signs of herpes infection include: cold sores, shingles, genital sores, skin rash, pain, and itching. Signs of more serious herpes infection include: changes in vision, eye redness or eye pain, severe or persistent headache, stiff neck, and confusion. Signs of infection can happen during treatment or after you have received your last dose of OCREVUS or OCREVUS ZUNOVO. Tell your healthcare provider right away if you have an infection. Your healthcare provider should delay your treatment with OCREVUS or OCREVUS ZUNOVO until your infection is gone.

- o **Hepatitis B virus (HBV) reactivation:** Before starting treatment with ocrelizumab, your healthcare provider will do blood tests to check for hepatitis B viral infection. If you have ever had hepatitis B virus infection, the hepatitis B virus may become active again during or after treatment with OCREVUS or OCREVUS ZUNOVO. Hepatitis B virus becoming active again (called reactivation) may cause serious liver problems including liver failure or death. Your healthcare provider will monitor you if you are at risk for hepatitis B virus reactivation during treatment and after you stop receiving OCREVUS or OCREVUS ZUNOVO.

- o **Weakened immune system:** OCREVUS or OCREVUS ZUNOVO taken before or after other medicines that weaken the immune system could increase your risk of getting infections.

- **Progressive Multifocal Leukoencephalopathy (PML):** PML is a rare brain infection that usually leads to death or severe disability and has been reported with ocrelizumab. Symptoms of PML get worse over days to weeks. It is important that you call your healthcare provider right away if you have any new or worsening neurologic signs or symptoms that have lasted several days, including problems with:

- o thinking
- o eyesight
- o strength
- o balance
- o weakness on 1 side of your body
- o using your arms or legs

- **Decreased immunoglobulins:** OCREVUS and OCREVUS ZUNOVO may cause a decrease in some types of immunoglobulins. Your healthcare provider will do blood tests to check your blood immunoglobulin levels.

Before receiving OCREVUS or OCREVUS ZUNOVO, tell your healthcare provider about all of your medical conditions, including if you:

- have or think you have an infection. See **“What is the most important information I should know about OCREVUS and OCREVUS ZUNOVO?”**
- have ever taken, take, or plan to take medicines that affect your immune system, or other treatments for MS. These medicines could increase your risk of getting an infection.

INDICATIONS AND IMPORTANT SAFETY INFORMATION (CONT.)

- have ever had hepatitis B or are a carrier of the hepatitis B virus.
- have a history of inflammatory bowel disease or colitis.
- have had a recent vaccination or are scheduled to receive any vaccinations.
 - **You should receive any required ‘live’ or ‘live-attenuated’ vaccines at least 4 weeks before you start treatment with OCREVUS or OCREVUS ZUNOVO.** You **should not receive** ‘live’ or ‘live-attenuated’ vaccines while you are being treated with OCREVUS or OCREVUS ZUNOVO and until your healthcare provider tells you that your immune system is no longer weakened.
 - **When possible, you should receive any ‘non-live’ vaccines at least 2 weeks before you start treatment with OCREVUS or OCREVUS ZUNOVO.** If you would like to receive any non-live (inactivated) vaccines, including the seasonal flu vaccine, while you are being treated with OCREVUS or OCREVUS ZUNOVO, talk to your healthcare provider.
 - If you have a baby and you received OCREVUS or OCREVUS ZUNOVO during your pregnancy, it is important to tell your baby’s healthcare provider about receiving OCREVUS or OCREVUS ZUNOVO so they can decide when your baby should be vaccinated.
- are pregnant, think that you might be pregnant, or plan to become pregnant. It is not known if OCREVUS and OCREVUS ZUNOVO will harm your unborn baby. You should use birth control (contraception) during treatment with OCREVUS and OCREVUS ZUNOVO and for 6 months after your last dose of OCREVUS or OCREVUS ZUNOVO. Talk with your healthcare provider about what birth control method is right for you during this time. Tell your healthcare provider if you become pregnant while receiving OCREVUS or OCREVUS ZUNOVO.
- are breastfeeding or plan to breastfeed. It is not known if OCREVUS and OCREVUS ZUNOVO pass into your breast milk. Talk to your healthcare provider about the best way to feed your baby if you take OCREVUS or OCREVUS ZUNOVO.

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

What are the possible side effects of OCREVUS and OCREVUS ZUNOVO?

OCREVUS and OCREVUS ZUNOVO may cause serious side effects, including:

- **Risk of cancers (malignancies) including breast cancer:** Follow your healthcare provider’s instructions about standard screening guidelines for breast cancer.
- **Inflammation of the colon, or colitis:** Tell your healthcare provider if you have any symptoms of colitis, such as:
 - Diarrhea (loose stools) or more frequent bowel movements than usual
 - Stools that are black, tarry, sticky or have blood or mucus
 - Severe stomach-area (abdomen) pain or tenderness

The most common side effects of OCREVUS ZUNOVO include:

- Injection reactions
- Respiratory tract infections
- Skin infections

INDICATIONS AND IMPORTANT SAFETY INFORMATION (CONT.)

These are not all the possible side effects of OCREVUS and OCREVUS ZUNOVO.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088. You may also report side effects to Genentech at (888) 835-2555.

For more information, go to www.OCREVUS.com or call 1-844-627-3887.

Please see additional Important Safety Information throughout and click here for full [Prescribing Information](#) and [Medication Guide for OCREVUS](#).

Please see additional Important Safety Information throughout and click here for full [Prescribing Information](#) and [Medication Guide for OCREVUS ZUNOVO](#).