

Treatment Referral Form

This is an optional form that can be used to provide information to your patient’s treatment site. **Please send this completed form to the treatment center that has been identified to administer OCREVUS® [IV] or OCREVUS ZUNOVO™ for your patient. Do not send it to Genentech.**

OCREVUS Patient Navigators are a point of contact for assistance throughout your patients’ treatment, including access, reimbursement and treatment coordination support. For more information, call (844) OCREVUS (844-627-3887).

1 PATIENT AND INSURANCE INFORMATION

Patient first name		Patient last name		Patient insurance		Please provide copies of the front and back of medical and pharmacy insurance cards.	
Address			City		Medical insurance		
State	ZIP	DOB (MM/DD/YYYY)		Insurance company name		Plan type	
		____ / ____ / ____					
Phone number				Member group number		ID number	
Preferred language, if not English				Policyholder name		Phone number	
Note: If possible, please provide MRI results and any supporting clinical notes, which include any past tried and/or failed therapies, intolerance, outcomes or contraindications to conventional therapy. Include any lab results and/or tests to support diagnosis.				Relationship to policyholder		Policyholder DOB (MM/DD/YYYY)	
		____ / ____ / ____					
Height	Weight	Temperature	Allergies	Pharmacy insurance			
Date of last MRI		Past DMT therapies		Prescription drug plan		Plan number	
Hepatitis B (HBsAg and anti-HBV) test results				Group number		ID number	
Quantitative serum immunoglobulins test results				Cardholder name		Phone number	
<input type="checkbox"/> Please confirm compliance: According to immunization guidelines, live or live-attenuated vaccines should be administered at least 4 weeks prior to initiation of OCREVUS or OCREVUS ZUNOVO and, whenever possible, for non-live vaccines at least 2 weeks prior to initiation of OCREVUS or OCREVUS ZUNOVO.				Relationship to cardholder		PCN/BIN number	

2 PRESCRIBER INFORMATION

Prescriber name		Prescriber NPI number		State license number	
Practice/facility name		Address		City	State
					ZIP
Primary contact name		Phone number		Fax number	

3

DIAGNOSIS/PRESCRIPTION INFORMATION

Please confirm diagnosis☐ G35 Multiple Sclerosis (MS)

☐ OCREVUS® [IV]

☐ Refills (# of refills): _____

☐ Dispense: 2 vials

Strength: 300 mg/10 mL (30 mg/mL) single-dose vial

Please select appropriate dosing and administration:

☐ Initial dose: 600-mg dose administered as 2 separate IV infusions 2 weeks apart

- First, infuse 300-mg IV over approximately 2.5 hours
- 2 weeks later, infuse 300-mg IV over approximately 2.5 hours

☐ Maintenance dose: 600-mg dose administered once every 24 weeks; choose from 2 infusion options:

☐ Option 1: Single infusion administered over approximately 3.5 to 4 hours

☐ Option 2: Single infusion administered over approximately 2 hours (for eligible patients who have not experienced a serious infusion reaction with any previous OCREVUS infusion)

Note: Observe the patient for at least 1 hour after the completion of the infusion. Infusions may be interrupted or slowed as needed. See the OCREVUS Prescribing Information for additional details.

OCREVUS premedications

☐ Methylprednisolone (or equivalent corticosteroid): 100 mg administered intravenously approximately 30 minutes prior to each OCREVUS infusion

☐ Antihistamine (e.g., diphenhydramine): Premedicate approximately 30 to 60 minutes prior to each OCREVUS infusion to further reduce the frequency and severity of infusion reactions

☐ Antipyretic (e.g., acetaminophen): The addition of an antipyretic may also be considered

☐ Other: _____

Infusion supplies

- Filter (0.2 or 0.22 micron in-line)
- Infusion-related reaction medications (i.e., Benadryl, epi-pen, etc.): _____
- Other: _____

☐ OCREVUS ZUNOVO™ [SUBCUTANEOUS INJECTION]

☐ Refills (# of refills): _____

☐ Dispense: 1 vial

Strength: 920 mg ocrelizumab and 23,000 units of hyaluronidase single-dose vial

Please select appropriate dosing and administration:

☐ 920 mg/23,000 units (920 mg ocrelizumab and 23,000 units of hyaluronidase)

- Administered subcutaneously in the abdomen approximately 10 minutes once every 24 weeks


OCREVUS ZUNOVO premedications

☐ Dexamethasone (or equivalent corticosteroid): 20 mg administered orally at least 30 minutes prior to administration

☐ Antihistamine (e.g., desloratadine): Administered orally at least 30 minutes prior to administration to reduce the risk of local and systemic injection reactions

☐ Antipyretic (e.g., acetaminophen): The addition of an antipyretic may also be considered

☐ Other: _____



Scan or click this QR code to download a detailed **Dosing and Administration Guide** for OCREVUS and OCREVUS ZUNOVO.

Injection supplies


- OCREVUS subcutaneous injection vial
- Syringe
- 21G stainless steel transfer needle
- Subcutaneous injection set (e.g., winged/butterfly) containing a 24-26G needle
- Syringe tip cap

Optional:

- Syringe pump

Note: For the initial dose, monitor the patient for at least 1 hour post-injection. For subsequent doses, monitor the patient for at least 15 minutes post-injection. See the OCREVUS ZUNOVO Prescribing Information for additional details.

Additional information and clinical notes:



Please scan or click this QR code to download an optional **Day-of-Treatment Checklist and Discharge Instructions** to return to the prescriber following the patient's treatment.

Please see the OCREVUS infusion or injection-related reaction protocol in the OCREVUS and OCREVUS ZUNOVO Prescribing Information. Please see additional Important Safety Information throughout and click here for full OCREVUS [Prescribing Information](#) and [Medication Guide](#). For OCREVUS ZUNOVO, click here for full [Prescribing Information](#) and [Medication Guide](#).

Genentech

A Member of the Roche Group

OCREVUS® is a registered trademark and OCREVUS ZUNOVO™ is a trademark of Genentech, Inc.

©2024 Genentech USA, Inc. So. San Francisco, CA All rights reserved. M-US-00004344(v3.0) 09/24

Page 2 of 2